

Table 1C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Demographic Indicators
Children Aged < 5 Years (2,3)

Total Records	Number
Individual Children	54,764
Total Records (All Visits)	58,076

Source of Data	Number	%
WIC	54,764	100.0
EPSDT	0	0.0
Child Health-MCH	0	0.0
Child Health-HMO	0	0.0
Headstart	0	0.0
School Health	0	0.0
Other	0	0.0
Total	54,764	100.0
<i>Excluded Unknown: 0 0.0 %</i>		

Race/Ethnic Distribution	Number	%
White, Not Hispanic	15,058	27.6
Black, Not Hispanic	13,998	25.7
Hispanic	23,940	43.9
American Indian/Alaskan Native	164	0.3
Asian/Pacific Islander	1,010	1.9
Multiple Races	339	0.6
All Other	0	0.0
Total	54,509	100.0
<i>Excluded Unknown: 255 0.5 %</i>		

Age Distribution (4)	Number	%
0 - 5 Months	15,027	27.4
6 - 11 Months	1,360	2.5
12 - 23 Months	12,494	22.8
24 - 35 Months	9,298	17.0
3 - 4 Years	16,585	30.3
Total	54,764	100.0

% of Poverty Level	Number	%
0-50	25,927	47.7
51-100	12,311	22.6
101-130	6,971	12.8
131-150	3,362	6.2
151-185	4,453	8.2
186-200	501	0.9
Over 200	844	1.6
Adjunctive Eligibility	0	0.0
Total	54,369	100.0
<i>Excluded Unknown 395 0.7%</i>		

Program Participation	Number	%
WIC	54,764	100.0
FoodStamps	54,764	21.6
Medicaid	54,764	84.7
TANF	54,764	10.9
<i>Excluded Unknown:</i>		
WIC	0	0.0
FoodStamps	0	0.0
Medicaid	0	0.0
TANF	0	0.0

Migrant Status	Number	%
Yes	14	*
No	53,368	100.0
Total	53,382	100.0
<i>Excluded Unknown 1,382 2.5%</i>		

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 2C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Health Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Health Indicators ⁽³⁾						
			Connecticut Prevalence		National Prevalence (Prior Year)	
			Number	%	Number	%
Birthweight (4)						
Low	< 2500 g		14,616	9.3	1,633,363	9.3
High	> 4000 g		14,616	7.0	1,633,363	6.8
Height and Weight						
Short Stature (5)	< 5th		53,180	4.0	6,600,078	6.4
	< 10th		53,180	8.0	6,600,078	11.4
Underweight (6)	< 5th		53,180	7.1	6,600,078	4.7
	< 10th		53,180	11.7	6,600,078	8.0
Overweight (6) ≥ 2 Yrs Overweight (6)	≥ 95th		53,180	12.7	6,600,078	14.1
	85th - < 95th		24,956	16.3	2,822,660	16.1
	≥ 95th		24,956	16.6	2,822,660	14.8
Anemia (7)						
Low Hb			35,927	8.8	3,744,224	14.2
Low Hct			1,105	6.0	506,123	9.0
Low Hb/Hct			37,032	8.7	4,020,684	13.7
Breastfeeding (8)						
Ever Breastfed			14,667	56.3	1,469,572	56.1
Breastfed At Least 6 Months			5,330	22.9	718,598	23.4
Breastfed At Least 12 Months			9,646	17.2	792,792	17.6
Smoking in Household			0	*	389,446	20.1
≤ 2 Hours TV Viewing/ Day (9)			0	*	64,128	40.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(9) Children aged 2 years and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 3C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Breastfeeding Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Breastfeeding Indicators ⁽³⁾				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Breastfeeding Duration ^(4,5)				
Ever Breastfed	14,667	56.3	1,469,572	56.1
Breastfed 1+ Weeks	11,211	61.0	1,074,221	52.1
Breastfed 2+ Weeks	6,914	58.6	862,442	49.1
Breastfed 3+ Weeks	4,843	57.2	714,332	46.7
Breastfed 4+ Weeks	3,737	54.7	626,309	44.5
Breastfed 6+ Weeks	2,826	47.0	536,440	39.6
Breastfed 2+ Months	2,604	38.4	509,857	35.1
Breastfed 3+ Months	2,949	30.9	549,816	30.0
Breastfed 4+ Months	3,668	28.2	623,490	27.0
Breastfed 5+ Months	4,494	25.8	688,285	24.9
Breastfed 6+ Months	5,330	22.9	718,598	23.4
Breastfed 9+ Months	7,562	19.0	752,987	19.0
Breastfed 12+ Months	9,646	17.2	792,792	17.6
Breastfed 18+ Months	4,932	11.6	595,234	12.2
Exclusive Breastfeeding ⁽⁶⁾				
Exclusively Breastfed at Least 3 Months	0	*	37,398	9.8
Exclusively Breastfed at Least 6 Months	0	*	39,388	6.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. For each breastfeeding duration category, the analysis includes only children who turned that age during the reporting period by/on their date of visit. For example, infants who turned 9 months old during the reporting period by/on their date of visit are included in the 9+ months analysis.

(5) It is possible the analysis will show that slightly more infants are breastfed at older ages, e.g. 1+ week of age, than at younger ages, e.g. at birth (ever breastfed), due to variation in infants included in each breastfeeding analysis category.

(6) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Comparison of Racial and Ethnic Distribution by Contributor
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Contributor								
Contributor	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Connecticut	54,764	27.5	25.6	43.7	0.3	1.8	0.6	0.5
Nation (Prior Year)	6,948,667	35.6	20.3	37.1	1.2	2.9	0.1	2.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Comparison of Age Distribution by Contributor
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution by Contributor				
Contributor	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Connecticut	54,764	29.9	22.8	47.3
Nation (Prior Year)	6,948,667	35.4	22.2	42.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3)												
Contributor	Birthweight (4)			Height and Weight						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Over- weight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th- <95th	% ≥95th	Number	%
Connecticut	14,616	9.3	7.0	53,180	4.0	7.1	12.7	24,956	16.3	16.6	37,032	8.7
Nation (Prior Year)	1,633,363	9.3	6.8	6,600,078	6.4	4.7	14.1	2,822,660	16.1	14.8	4,020,684	13.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor
(4)

Contributor	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	%	Number	%		
Connecticut	14,667	56.3	5,330	22.9	9,646	17.2	0	*	0	*	0	*		
Nation (Prior Year)	1,469,572	56.1	718,598	23.4	792,792	17.6	37,398	9.8	39,388	6.7	64,128	40.5	389,446	20.1

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(6) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(7) Children ages 2 years and older included in the analyses.

Table 8C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Growth and Anemia Indicators by Race/Ethnicity or Age
Children Aged < 5 Years ⁽²⁾

Growth and Anemia Indicators by Race/Ethnicity or Age ⁽³⁾												
	Birthweight (4)			Height and Weight						Anemia Low Hb/Hct (7)		
	Number	Low %	High %	Number	Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		<2500 g	>4000 g		<5th	<5th	≥95th	Number	% 85th-<95th	% ≥95th	Number	%
Race/Ethnicity												
White, Not Hispanic	4,217	8.9	8.6	14,684	4.4	7.7	11.0	6,542	15.5	14.2	9,856	6.7
Black, Not Hispanic	3,565	11.8	5.4	13,491	3.6	8.0	11.4	6,422	14.4	14.3	9,656	11.9
Hispanic	6,342	8.1	6.8	23,289	3.8	6.1	14.9	11,208	17.9	19.9	16,332	8.1
American Indian/Alaskan Native	49	*	*	157	5.1	12.1	10.2	69	*	*	102	7.8
Asian/Pacific Islander	286	10.5	4.9	983	5.3	10.8	7.7	456	13.2	8.3	689	8.1
Multiple Races	131	7.6	10.7	327	4.0	8.0	9.8	129	17.1	9.3	190	9.5
All Other	26	*	*	249	4.0	4.8	14.1	130	19.2	13.8	207	8.2
Total	14,616	9.3	7.0	53,180	4.0	7.1	12.7	24,956	16.3	16.6	37,032	8.7
Age												
0 - 11 Months	-	-	-	16,254	4.1	12.1	5.6	-	*	*	2,566	10.4
12 - 23 Months	-	-	-	11,970	4.7	5.7	14.3	-	*	*	11,518	9.9
24 - 35 Months	-	-	-	9,106	3.2	6.2	13.0	9,106	14.9	13.0	8,751	9.2
36 - 47 Months	-	-	-	7,744	3.9	3.8	18.1	7,744	17.2	18.1	7,598	7.7
48 - 59 Months	-	-	-	8,106	3.5	3.5	19.4	8,106	17.0	19.4	6,599	6.7
Total	-	-	-	53,180	4.0	7.1	12.7	24,956	16.3	16.6	37,032	8.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 9C
2005 Pediatric Nutrition Surveillance
Connecticut

Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity or Age (2)
Children Aged < 5 Years (3)

Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity or Age (4)										
	Breastfeeding (5)						<2 Hours TV/Video Viewing/Day (6)		Smoking in Household	
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	4,232	52.9	1,501	21.7	2,710	16.6	0	*	0	*
Black, Not Hispanic	3,590	49.4	1,362	17.8	2,506	14.2	0	*	0	*
Hispanic	6,352	61.9	2,311	26.1	4,133	18.9	0	*	0	*
American Indian/Alaskan Native	50	*	15	*	28	*	0	*	0	*
Asian/Pacific Islander	286	68.5	104	31.7	173	23.7	0	*	0	*
Multiple Races	131	54.2	31	*	60	*	0	*	0	*
All Other	26	*	6	*	36	*	0	*	0	*
Total	14,667	56.3	5,330	22.9	9,646	17.2	0	*	0	*
Age										
0 - 11 Months	-	-	-	-	-	-	0	*	0	*
12 - 23 Months	-	-	-	-	-	-	0	*	0	*
24 - 35 Months	-	-	-	-	-	-	0	*	0	*
36 - 47 Months	-	-	-	-	-	-	0	*	0	*
48 - 59 Months	-	-	-	-	-	-	0	*	0	*
Total	14,667	56.3	5,330	22.9	9,646	17.2	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 10C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Racial and Ethnic Distribution
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Racial and Ethnic Distribution								
Year	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
2005	54,764	27.5	25.6	43.7	0.3	1.8	0.6	0.5
2004	56,050	27.8	26.5	41.4	0.2	1.5	0.2	2.3
2003	46,886	28.0	26.5	40.4	0.2	1.3	0.0	3.7
2002	761	31.1	24.6	38.0	0.3	1.6	0.0	4.5
2001	51,410	29.6	27.7	38.4	0.2	1.2	0.0	2.9
2000	857	35.8	28.5	32.1	0.0	0.8	0.0	2.8
1999	13	*	*	*	*	*	*	*
1998	53,177	31.8	29.7	35.2	0.2	1.2	0.0	2.0
1997	49,384	32.9	29.3	34.7	0.2	1.3	0.0	1.6
1996	0	*	*	*	*	*	*	*
1995	5,838	36.4	25.9	36.2	1.2	0.3	0.0	0.0
1994	6,823	38.8	26.2	33.8	1.1	0.2	0.0	0.0
1993	6,625	39.3	26.2	33.3	1.0	0.3	0.0	0.0
1992	3,587	37.9	29.5	31.7	0.8	0.2	0.0	0.0
1991	6,882	42.9	26.0	30.3	0.6	0.1	0.0	0.0
1990	6,002	42.8	28.4	28.1	0.5	0.1	0.0	0.0
1989	5,867	44.7	28.7	25.6	0.7	0.2	0.0	0.0
1988	5,907	51.0	26.3	21.7	0.6	0.4	0.0	0.0
1987	5,234	54.2	25.1	19.9	0.6	0.3	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 10C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Racial and Ethnic Distribution
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Racial and Ethnic Distribution								
Year	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1986	2,396	59.2	28.1	11.7	0.7	0.3	0.0	0.0
1985	2,344	65.9	23.6	9.4	0.8	0.4	0.0	0.0
1984	2,529	70.1	19.4	9.6	0.6	0.4	0.0	0.0
1983	2,009	77.1	11.1	10.0	0.4	1.3	0.0	0.0
1982	3,232	57.3	17.8	23.2	0.3	1.4	0.0	0.0
1981	40	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 11C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Trends in Age Distribution
Children Aged < 5 Years (2)

Summary of Trends in Age Distribution (3)				
Year	Number Included in Analysis (4)	0-11 Months %	12-23 Months %	24-59 Months %
2005	54,764	29.9	22.8	47.3
2004	56,050	28.8	22.8	48.5
2003	46,886	27.9	22.7	49.4
2002	761	33.1	26.0	40.9
2001	51,410	34.7	21.1	44.1
2000	857	45.6	15.5	38.9
1999	13	*	*	*
1998	53,177	34.8	20.4	44.8
1997	49,384	26.8	22.5	50.7
1996	0	*	*	*
1995	5,838	34.8	18.6	46.6
1994	6,823	34.7	18.3	47.0
1993	6,625	33.8	18.9	47.4
1992	3,587	33.0	20.2	46.8
1991	6,882	35.3	20.0	44.6
1990	6,002	36.2	19.5	44.3
1989	5,867	35.0	19.6	45.4
1988	5,907	35.0	20.2	44.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Records rejected if date of birth and date of visit unknown.

(4) Excludes records with unknown data or errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 11C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Trends in Age Distribution
Children Aged < 5 Years (2)

Summary of Trends in Age Distribution (3)				
Year	Number Included in Analysis (4)	0-11 Months %	12-23 Months %	24-59 Months %
1987	5,234	33.0	19.3	47.7
1986	2,396	37.1	22.3	40.7
1985	2,344	42.0	21.2	36.7
1984	2,529	53.6	16.5	29.9
1983	2,009	59.6	15.6	24.8
1982	3,232	59.8	13.6	26.6
1981	40	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Records rejected if date of birth and date of visit unknown.

(4) Excludes records with unknown data or errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Number	%	
	Number	% <2500 g	% >4000 g									Number
2005	14,616	9.3	7.0	53,180	4.0	7.1	12.7	24,956	16.3	16.6	37,032	8.7
2004	14,430	9.2	6.9	54,429	4.0	7.0	13.1	26,109	15.9	16.9	38,724	9.4
2003	10,899	9.7	6.9	45,457	6.0	5.1	16.8	22,495	17.6	19.6	36,004	9.9
2002	215	10.7	7.0	649	4.8	7.9	14.3	280	18.2	19.6	554	10.1
2001	13,868	9.0	7.9	50,032	6.2	5.2	16.3	22,083	17.3	18.7	39,670	10.4
2000	376	7.4	9.0	829	3.3	5.9	13.6	319	17.9	16.6	591	10.7
1999	0	*	*	2	*	*	*	0	*	*	5	*
1998	14,129	10.3	8.0	51,780	6.7	5.3	15.4	23,272	17.8	17.8	41,153	10.8
1997	8,817	9.4	8.1	48,370	6.9	4.9	15.9	24,511	17.1	16.6	42,707	10.8
1996	0	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1995	992	8.9	8.9	5,683	4.7	6.6	11.5	2,659	15.0	13.2	3,407	9.4
1994	1,448	7.9	8.6	6,620	4.8	7.4	11.2	3,096	13.8	12.7	4,269	7.3
1993	1,215	10.1	6.0	6,431	4.7	7.4	10.7	3,068	12.9	11.8	4,065	6.0
1992	623	8.2	8.7	3,469	5.2	7.5	12.2	1,639	13.8	12.6	1,911	8.7
1991	1,401	8.3	7.6	6,582	4.6	8.2	10.5	2,952	14.7	11.3	4,497	9.4
1990	1,204	9.1	8.0	5,705	5.0	7.4	10.5	2,588	14.5	11.0	3,907	7.8
1989	1,237	10.3	7.0	5,570	4.8	6.8	10.4	2,582	13.7	11.0	3,909	8.4
1988	1,092	7.9	9.2	5,598	4.5	7.5	10.1	2,556	12.9	11.0	3,886	8.6
1987	1,039	8.3	9.9	4,983	4.7	7.2	9.3	2,416	13.6	9.9	3,209	7.5
1986	487	8.4	9.2	2,320	5.9	6.4	10.6	949	12.8	8.7	1,498	7.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under-weight (6)	Over-weight (6)	≥ 2 Yrs Overweight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1985	615	9.1	7.8	2,268	6.7	8.1	8.8	852	11.0	6.7	1,506	11.0
1984	926	9.4	9.9	2,420	6.2	9.1	6.4	723	12.3	6.6	1,311	10.1
1983	844	8.9	8.8	1,928	6.6	12.6	7.6	488	8.8	7.2	978	9.6
1982	1,653	9.7	6.2	2,991	8.5	11.3	6.4	838	13.0	7.6	1,371	10.9
1981	30	*	*	26	*	*	*	4	*	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 13C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Breastfeeding
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Breastfeeding ⁽³⁾						
Year	Breastfeeding ⁽⁴⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
2005	14,667	56.3	5,330	22.9	9,646	17.2
2004	14,459	55.3	5,725	22.9	10,005	17.2
2003	10,899	53.8	4,793	23.0	7,741	16.5
2002	215	56.7	151	23.8	183	15.3
2001	13,868	49.3	8,784	23.4	7,977	17.7
2000	377	46.4	196	15.8	131	11.5
1999	0	*	7	*	1	*
1998	14,129	42.7	8,760	22.4	7,588	16.8
1997	8,817	40.2	8,818	17.5	7,760	11.8
1996	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Breastfeeding
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Breastfeeding ⁽³⁾						
Year	Breastfeeding ⁽⁴⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
1995	788	35.7	300	11.3	197	7.1
1994	1,151	33.7	628	12.4	447	5.1
1993	995	33.3	508	9.6	378	4.5
1992	556	26.1	308	8.1	237	4.2
1991	1,286	25.8	814	7.4	689	3.6
1990	1,087	29.8	665	8.3	561	2.5
1989	1,152	26.1	788	7.0	645	3.7
1988	1,047	26.7	837	6.6	618	2.6
1987	1,002	25.1	736	7.3	613	4.4
1986	487	37.4	354	12.4	298	3.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Summary of Trends in Breastfeeding
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Breastfeeding ⁽³⁾						
	Breastfeeding ⁽⁴⁾					
Year	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
1985	613	36.4	458	7.2	344	4.9
1984	925	38.9	479	15.0	332	8.1
1983	845	40.4	347	15.9	191	11.0
1982	1,653	30.7	370	13.0	245	4.9
1981	30	*	0	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 14C
2005 Pediatric Nutrition Surveillance
Connecticut
Children Aged < 5 Years

Page: 1
Run Date: 2 / 2 / 07

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 15C
2005 Pediatric Nutrition Surveillance
Connecticut
Children Aged < 5 Years

Page: 1
Run Date: 2 / 2 / 07

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 16C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Growth Indicators by Race/Ethnicity and Age
Children Aged < 5 Years ⁽²⁾

Growth Indicators by Race/Ethnicity and Age ⁽³⁾									
	Short Stature (4)		Underweight (5)		Overweight (5)		≥ 2 Yrs Overweight (5)		
	Number	% <5th	Number	% <5th	Number	% ≥95th	Number	% 85th - <95th	% ≥95th
White, Not Hispanic									
0 - 11 Months	4,771	4.5	4,771	12.7	4,771	5.4	0	0.0	0.0
12 - 23 Months	3,371	4.5	3,371	6.3	3,371	12.7	0	0.0	0.0
24 - 59 Months	6,542	4.3	6,542	4.8	6,542	14.2	6,542	15.5	14.2
Total	14,684	4.4	14,684	7.7	14,684	11.0	6,542	15.5	14.2
Black, Not Hispanic									
0 - 11 Months	3,955	4.7	3,955	13.5	3,955	5.3	0	0.0	0.0
12 - 23 Months	3,114	4.7	3,114	6.5	3,114	13.1	0	0.0	0.0
24 - 59 Months	6,422	2.4	6,422	5.4	6,422	14.3	6,422	14.4	14.3
Total	13,491	3.6	13,491	8.0	13,491	11.4	6,422	14.4	14.3
Hispanic									
0 - 11 Months	6,973	3.5	6,973	10.8	6,973	5.8	0	0.0	0.0
12 - 23 Months	5,108	4.7	5,108	4.6	5,108	16.3	0	0.0	0.0
24 - 59 Months	11,208	3.7	11,208	3.8	11,208	19.9	11,208	17.9	19.9
Total	23,289	3.8	23,289	6.1	23,289	14.9	11,208	17.9	19.9
American Indian/Alaskan Native									
0 - 11 Months	51	*	51	*	51	*	0	*	*
12 - 23 Months	37	*	37	*	37	*	0	*	*
24 - 59 Months	69	*	69	*	69	*	69	*	*
Total	157	5.1	157	12.1	157	10.2	69	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 16C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Connecticut
Growth Indicators by Race/Ethnicity and Age
Children Aged < 5 Years ⁽²⁾

Growth Indicators by Race/Ethnicity and Age ⁽³⁾									
	Short Stature (4)		Underweight (5)		Overweight (5)		≥ 2 Yrs Overweight (5)		
	Number	% <5th	Number	% <5th	Number	% ≥95th	Number	% 85th - <95th	% ≥95th
Asian/Pacific Islander									
0 - 11 Months	316	4.7	316	15.2	316	5.4	0	0.0	0.0
12 - 23 Months	211	8.5	211	10.0	211	10.0	0	0.0	0.0
24 - 59 Months	456	4.2	456	8.1	456	8.3	456	13.2	8.3
Total	983	5.3	983	10.8	983	7.7	456	13.2	8.3
Multiple Races									
0 - 11 Months	135	3.7	135	10.4	135	8.9	0	0.0	0.0
12 - 23 Months	63	*	63	*	63	*	0	*	*
24 - 59 Months	129	3.9	129	7.0	129	9.3	129	17.1	9.3
Total	327	4.0	327	8.0	327	9.8	129	17.1	9.3
All Other/Unknown									
0 - 11 Months	53	*	53	*	53	*	0	*	*
12 - 23 Months	66	*	66	*	66	*	0	*	*
24 - 59 Months	130	4.6	130	3.8	130	13.8	130	19.2	13.8
Total	249	4.0	249	4.8	249	14.1	130	19.2	13.8
Total All Race/Ethnic Groups									
0 - 11 Months	16,254	4.1	16,254	12.1	16,254	5.6	0	0.0	0.0
12 - 23 Months	11,970	4.7	11,970	5.7	11,970	14.3	0	0.0	0.0
24 - 59 Months	24,956	3.5	24,956	4.6	24,956	16.6	24,956	16.3	16.6
Total	53,180	4.0	53,180	7.1	53,180	12.7	24,956	16.3	16.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Anemia Indicators by Race/Ethnicity and Age
Children Aged < 5 Years (2)

Page: 1
Run Date: 2 / 2 / 07

Anemia Indicators by Race/Ethnicity and Age (3, 4)					
	Anemia Low Hb/Hct %			Anemia Low Hb/Hct %	
	Number		Number		
White, Not Hispanic			Asian/Pacific Islander		
6 - 11 Months	794	7.7	6 - 11 Months	51	*
12 - 17 Months	1,922	7.4	12 - 17 Months	132	7.6
18 - 23 Months	1,263	7.5	18 - 23 Months	76	*
24 - 35 Months	2,276	5.8	24 - 35 Months	151	8.6
36 - 59 Months	3,601	6.5	36 - 59 Months	279	6.5
Total	9,856	6.7	Total	689	8.1
Black, Not Hispanic			Multiple Races		
6 - 11 Months	632	11.6	6 - 11 Months	14	*
12 - 17 Months	1,880	11.9	12 - 17 Months	35	*
18 - 23 Months	1,131	14.4	18 - 23 Months	25	*
24 - 35 Months	2,261	13.7	24 - 35 Months	60	*
36 - 59 Months	3,752	10.2	36 - 59 Months	56	*
Total	9,656	11.9	Total	190	9.5
Hispanic			All Other/Unknown		
6 - 11 Months	1,040	11.7	6 - 11 Months	20	*
12 - 17 Months	2,969	9.7	12 - 17 Months	41	*
18 - 23 Months	1,993	9.8	18 - 23 Months	23	*
24 - 35 Months	3,944	8.6	24 - 35 Months	34	*
36 - 59 Months	6,386	6.0	36 - 59 Months	89	*
Total	16,332	8.1	Total	207	8.2
American Indian/Alaskan Native			Total All Race/Ethnic Groups		
6 - 11 Months	15	*	6 - 11 Months	2,566	10.4
12 - 17 Months	17	*	12 - 17 Months	6,996	9.6
18 - 23 Months	11	*	18 - 23 Months	4,522	10.3
24 - 35 Months	25	*	24 - 35 Months	8,751	9.2
36 - 59 Months	34	*	36 - 59 Months	14,197	7.2
Total	102	7.8	Total	37,032	8.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th		
White, Not Hispanic	2005	4,217	8.9	8.6	14,684	4.4	7.7	11.0	6,542	15.5	14.2	9,856	6.7
	2004	4,283	8.6	9.0	15,194	4.5	7.4	10.9	6,887	15.5	14.4	10,335	7.1
	2003	3,321	8.8	8.9	12,755	6.3	5.2	14.4	6,067	18.5	16.2	9,739	7.4
	2002	77	*	*	212	5.2	6.6	17.9	84	*	*	165	6.7
	2001	4,325	8.4	9.9	14,869	6.4	5.6	13.7	6,259	17.5	15.2	11,413	7.7
	2000	150	7.3	11.3	295	4.1	5.4	11.9	98	*	*	195	6.7
	1999	0	*	*	0	*	*	*	0	*	*	0	*
	1998	4,918	9.2	10.4	16,507	6.5	6.1	12.8	6,861	16.4	14.8	12,573	7.4
	1997	3,392	8.1	10.4	15,980	7.1	5.4	13.3	7,397	16.8	13.9	13,452	7.8
Black, Not Hispanic	2005	3,565	11.8	5.4	13,491	3.6	8.0	11.4	6,422	14.4	14.3	9,656	11.9
	2004	3,829	11.9	5.4	14,373	3.7	7.8	11.7	7,009	15.5	14.4	10,361	12.9
	2003	2,848	12.1	5.6	11,993	5.8	5.8	15.7	5,905	16.2	17.3	9,590	13.4
	2002	50	*	*	154	3.9	12.3	7.8	66	*	*	140	13.6
	2001	3,815	11.5	6.2	13,810	6.3	5.5	15.7	6,132	16.7	16.5	11,119	14.3
	2000	105	7.6	6.7	236	2.1	7.2	14.4	91	*	*	174	13.8
	1999	0	*	*	0	*	*	*	0	*	*	1	*
	1998	4,194	12.6	6.6	15,322	6.4	5.6	14.8	6,904	17.2	16.0	12,333	14.8
	1997	2,635	11.1	5.7	14,119	6.5	5.4	15.4	7,166	16.1	15.3	12,584	14.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)					
		Number	% <2500 g	% >4000 g				Number	% <5th	% <5th	% ≥95th		
Hispanic	2005	6,342	8.1	6.8	23,289	3.8	6.1	14.9	11,208	17.9	19.9	16,332	8.1
	2004	5,669	7.9	6.4	22,520	3.8	5.9	15.6	11,147	16.4	20.1	16,404	8.7
	2003	4,107	9.0	6.4	18,364	6.0	4.1	19.6	9,478	18.2	23.3	14,887	9.1
	2002	71	*	*	242	5.4	5.8	15.7	119	20.2	20.2	220	10.9
	2001	5,078	7.5	7.6	19,204	6.0	4.5	19.2	8,849	17.7	23.1	15,470	9.7
	2000	100	6.0	9.0	267	3.7	5.6	15.0	121	20.7	21.5	206	12.6
	1999	0	*	*	2	*	*	*	0	*	*	3	*
	1998	4,504	9.7	6.7	18,204	7.1	4.3	18.4	8,815	19.5	21.9	14,897	10.2
	1997	2,491	8.9	7.1	16,771	7.1	3.6	19.2	9,282	18.4	19.9	15,380	10.4
American Indian/Alaskan Native	2005	49	*	*	157	5.1	12.1	10.2	69	*	*	102	7.8
	2004	34	*	*	117	5.1	7.7	9.4	57	*	*	85	*
	2003	19	*	*	89	*	*	*	44	*	*	70	*
	2002	1	*	*	2	*	*	*	0	*	*	2	*
	2001	29	*	*	90	*	*	*	36	*	*	70	*
	2000	0	*	*	0	*	*	*	0	*	*	0	*
	1999	0	*	*	0	*	*	*	0	*	*	0	*
	1998	33	*	*	98	*	*	*	46	*	*	74	*
	1997	14	*	*	119	3.4	7.6	9.2	81	*	*	111	9.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)					
		Number	% <2500 g	% >4000 g				Number	% <5th	% <5th	% ≥95th		
Asian/Pacific Islander	2005	286	10.5	4.9	983	5.3	10.8	7.7	456	13.2	8.3	689	8.1
	2004	254	12.2	7.1	843	3.9	12.2	10.3	367	9.5	14.7	568	7.7
	2003	149	8.7	2.7	575	6.3	7.1	13.4	270	13.0	17.8	435	9.0
	2002	4	*	*	11	*	*	*	4	*	*	8	*
	2001	217	10.6	6.0	608	6.7	8.7	12.7	222	14.0	16.7	446	9.6
	2000	5	*	*	7	*	*	*	2	*	*	3	*
	1999	0	*	*	0	*	*	*	0	*	*	0	*
	1998	173	8.1	4.6	622	6.9	7.9	13.8	268	15.7	16.4	482	9.8
	1997	127	11.8	6.3	624	4.8	7.1	11.5	299	14.4	13.4	532	9.6
Multiple Races	2005	131	7.6	10.7	327	4.0	8.0	9.8	129	17.1	9.3	190	9.5
	2004	35	*	*	119	0.8	8.4	7.6	43	*	*	82	*
	2003	0	*	*	0	*	*	*	0	*	*	0	*
	2002	0	*	*	0	*	*	*	0	*	*	0	*
	2001	0	*	*	0	*	*	*	0	*	*	0	*
	2000	0	*	*	0	*	*	*	0	*	*	0	*
	1999	0	*	*	0	*	*	*	0	*	*	0	*
	1998	0	*	*	0	*	*	*	0	*	*	0	*
	1997	0	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)		Under-weight (6)	Over-weight (6)	≥ 2 Yrs Overweight (6)				
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th		
All Other/Unknown	2005	26	*	*	249	4.0	4.8	14.1	130	19.2	13.8	207	8.2
	2004	326	7.4	6.4	1,263	4.0	9.4	13.1	599	17.0	16.4	889	8.1
	2003	455	8.1	6.6	1,681	5.3	7.6	14.6	731	16.1	17.4	1,283	11.5
	2002	12	*	*	28	*	*	*	7	*	*	19	*
	2001	404	9.4	7.7	1,451	5.7	7.0	12.3	585	17.4	13.2	1,152	10.9
	2000	16	*	*	24	*	*	*	7	*	*	13	*
	1999	0	*	*	0	*	*	*	0	*	*	1	*
	1998	307	7.2	8.5	1,027	6.1	5.5	13.6	378	17.5	15.1	794	12.5
	1997	158	14.6	13.3	757	8.1	8.3	14.3	286	14.0	12.9	648	9.7
Total All Race/Ethnic Groups	2005	14,616	9.3	7.0	53,180	4.0	7.1	12.7	24,956	16.3	16.6	37,032	8.7
	2004	14,430	9.2	6.9	54,429	4.0	7.0	13.1	26,109	15.9	16.9	38,724	9.4
	2003	10,899	9.7	6.9	45,457	6.0	5.1	16.8	22,495	17.6	19.6	36,004	9.9
	2002	215	10.7	7.0	649	4.8	7.9	14.3	280	18.2	19.6	554	10.1
	2001	13,868	9.0	7.9	50,032	6.2	5.2	16.3	22,083	17.3	18.7	39,670	10.4
	2000	376	7.4	9.0	829	3.3	5.9	13.6	319	17.9	16.6	591	10.7
	1999	0	*	*	2	*	*	*	0	*	*	5	*
	1998	14,129	10.3	8.0	51,780	6.7	5.3	15.4	23,272	17.8	17.8	41,153	10.8
	1997	8,817	9.4	8.1	48,370	6.9	4.9	15.9	24,511	17.1	16.6	42,707	10.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
White, Not Hispanic	2005	4,232	52.9	1,501	21.7	2,710	16.6
	2004	4,288	51.5	1,645	20.9	2,834	14.4
	2003	3,321	48.9	1,257	18.1	2,152	13.8
	2002	77	*	47	*	59	*
	2001	4,325	44.7	2,510	20.2	2,368	15.7
	2000	150	48.0	70	*	48	*
	1999	0	*	4	*	0	*
	1998	4,918	43.1	2,785	21.9	2,306	16.5
	1997	3,392	42.5	2,961	17.3	2,536	12.7
Black, Not Hispanic	2005	3,590	49.4	1,362	17.8	2,506	14.2
	2004	3,840	49.5	1,523	20.3	2,597	14.9
	2003	2,848	49.8	1,315	20.5	2,064	15.7
	2002	50	*	35	*	47	*
	2001	3,815	43.0	2,551	23.1	2,150	16.8
	2000	105	45.7	57	*	40	*
	1999	0	*	0	*	0	*
	1998	4,194	37.9	2,777	20.1	2,273	15.0
	1997	2,635	34.6	2,647	15.9	2,286	9.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
Hispanic	2005	6,352	61.9	2,311	26.1	4,133	18.9
	2004	5,680	60.7	2,331	25.4	4,170	20.0
	2003	4,107	58.6	1,904	26.6	3,090	17.6
	2002	71	*	59	*	62	*
	2001	5,078	56.4	3,296	24.6	3,084	18.3
	2000	101	45.5	62	*	42	*
	1999	0	*	1	*	1	*
	1998	4,504	45.4	2,870	24.4	2,753	17.6
	1997	2,491	42.0	2,889	18.7	2,679	12.0
American Indian/Alaskan Native	2005	50	*	15	*	28	*
	2004	34	*	12	*	21	*
	2003	19	*	11	*	17	*
	2002	1	*	1	*	1	*
	2001	29	*	13	*	15	*
	2000	0	*	0	*	0	*
	1999	0	*	0	*	0	*
	1998	33	*	20	*	12	*
	1997	14	*	9	*	11	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 19C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
Asian/Pacific Islander	2005	286	68.5	104	31.7	173	23.7
	2004	255	63.5	90	*	158	23.4
	2003	149	59.1	78	*	98	*
	2002	4	*	2	*	4	*
	2001	217	48.8	125	18.4	86	*
	2000	5	*	1	*	0	*
	1999	0	*	0	*	0	*
	1998	173	44.5	102	28.4	89	*
	1997	127	38.6	137	21.9	93	*
Multiple Races	2005	131	54.2	31	*	60	*
	2004	35	*	12	*	33	*
	2003	0	*	0	*	0	*
	2002	0	*	0	*	0	*
	2001	0	*	0	*	0	*
	2000	0	*	0	*	0	*
	1999	0	*	0	*	0	*
	1998	0	*	0	*	0	*
	1997	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 19C
2005 Pediatric Nutrition Surveillance (1)
Connecticut

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
All Other/Unknown	2005	26	*	6	*	36	*
	2004	327	69.1	112	27.7	192	25.0
	2003	455	70.8	228	33.3	320	28.1
	2002	12	*	7	*	10	*
	2001	404	69.1	289	41.2	274	31.8
	2000	16	*	6	*	1	*
	1999	0	*	2	*	0	*
	1998	307	58.3	206	30.6	155	21.3
	1997	158	57.0	175	21.7	155	20.6
Total All Race/Ethnic Groups	2005	14,667	56.3	5,330	22.9	9,646	17.2
	2004	14,459	55.3	5,725	22.9	10,005	17.2
	2003	10,899	53.8	4,793	23.0	7,741	16.5
	2002	215	56.7	151	23.8	183	15.3
	2001	13,868	49.3	8,784	23.4	7,977	17.7
	2000	377	46.4	196	15.8	131	11.5
	1999	0	*	7	*	1	*
	1998	14,129	42.7	8,760	22.4	7,588	16.8
	1997	8,817	40.2	8,818	17.5	7,760	11.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 20C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Trends in Growth Indicators by Age
Children Aged < 5 Years (2)

Summary of Trends in Growth Indicators by Age (3)																					
Age and Year *	Height and Weight								Height and Weight												
	Short Stature (4)	Under- weight (5)	Over- weight (5)	≥ 2 Yrs Overweight (5)					Short Stature (4)	Under- weight (5)	Over- weight (5)	≥ 2 Yrs Overweight (5)									
	Number	%	%	%	Number				%	%	Number	%	%	%	Number	%	%				
	<5th	<5th	≥95th	85th-<95th	≥95th			<5th	<5th	≥95th	85th-<95th	≥95th			<5th	<5th	≥95th	85th-<95th	≥95th		
0 - 11 Months																					
2005	16,254	4.1	12.1	5.6	-	-	-														
2004	15,989	4.3	12.2	6.0	-	-	-														
2003	12,437	6.8	8.4	11.6	-	-	-														
2002	209	4.8	9.6	10.0	-	-	-														
2001	17,237	6.8	7.5	11.7	-	-	-														
2000	378	2.9	6.6	10.1	-	-	-														
1999	1	*	*	*	-	-	-														
1998	17,801	6.8	8.1	10.8	-	-	-														
1997	12,888	7.9	7.1	12.8	-	-	-														
					0	-	-*														
12 - 23 Months																					
2005	11,970	4.7	5.7	14.3	-	-	-														
2004	12,331	4.6	6.0	14.2	-	-	-														
2003	10,525	7.3	4.4	17.3	-	-	-														
2002	160	5.0	10.6	10.6	-	-	-														
2001	10,712	8.2	4.2	18.9	-	-	-														
2000	132	4.5	7.6	16.7	-	-	-														
1999	1	*	*	*	-	-	-														
1998	10,707	7.8	4.4	17.6	-	-	-														
1997	10,971	8.6	4.4	18.2	-	-	-														
					0	-	-*														
24 - 35 Months																					
2005	9,106	3.2	6.2	13.0	9,106	14.9	13.0														
2004	9,346	3.3	5.6	14.5	9,346	13.7	14.5														
2003	8,380	4.8	4.4	17.1	8,380	16.3	17.1														
2002	74	*	*	*	74	*	*														
2001	8,294	4.8	5.1	16.4	8,294	15.7	16.4														
2000	93	*	*	*	93	*	*														
1999	0	*	*	*	0	*	*														
1998	8,408	6.0	4.4	16.3	8,408	16.2	16.3														
1997	8,676	5.7	4.4	15.0	8,676	16.3	15.0														
					0	*	*														
36 - 47 Months																					
2005	7,744	3.9	3.8	18.1	7,744	17.2	18.1														
2004	8,209	3.7	3.8	18.5	8,209	16.7	18.5														
2003	6,906	6.1	3.1	21.5	6,906	17.8	21.5														
2002	66	*	*	*	66	*	*														
2001	6,970	5.1	3.4	19.7	6,970	18.7	19.7														
2000	72	*	*	*	72	*	*														
1999	0	*	*	*	0	*	*														
1998	7,408	6.5	3.4	18.9	7,408	18.4	18.9														
1997	7,712	5.7	3.4	17.7	7,712	17.3	17.7														
					0	*	*														

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 20C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Trends in Growth Indicators by Age
Children Aged < 5 Years (2)

Summary of Trends in Growth Indicators by Age (3)																		
Age and Year *	Height and Weight					Age and Year *	Height and Weight											
	Short Stature (4)	Under- weight (5)	Over- weight (5)	≥ 2 Yrs Overweight (5)			Short Stature (4)	Under- weight (5)	Over- weight (5)	≥ 2 Yrs Overweight (5)								
	Number	%	%	%	Number		%	%	Number	%	%	Number	%	%				
	<5th	<5th	≥95th	85th-<95th	≥95th		<5th	<5th	≥95th	85th-<95th	≥95th		<5th	<5th	≥95th	85th-<95th	≥95th	
48 - 59 Months																		
2005	8,106	3.5	3.5	19.4	8,106	17.0	19.4											
2004	8,554	3.4	3.4	17.9	8,554	17.5	17.9											
2003	7,209	4.2	2.9	20.5	7,209	18.9	20.5											
2002	140	5.0	3.6	20.0	140	23.6	20.0											
2001	6,819	4.5	2.9	20.4	6,819	17.9	20.4											
2000	154	2.6	3.9	16.2	154	18.2	16.2											
1999	0	*	*	*	0	*	*											
1998	7,456	5.5	2.9	18.5	7,456	19.1	18.5											
1997	8,123	5.4	3.6	17.2	8,123	17.8	17.2											
					0	*	*											
Total All Age Groups																		
2005	53,180	4.0	7.1	12.7	24,956	16.3	16.6											
2004	54,429	4.0	7.0	13.1	26,109	15.9	16.9											
2003	45,457	6.0	5.1	16.8	22,495	17.6	19.6											
2002	649	4.8	7.9	14.3	280	18.2	19.6											
2001	50,032	6.2	5.2	16.3	22,083	17.3	18.7											
2000	829	3.3	5.9	13.6	319	17.9	16.6											
1999	2	*	*	*	0	*	*											
1998	51,780	6.7	5.3	15.4	23,272	17.8	17.8											
1997	48,370	6.9	4.9	15.9	24,511	17.1	16.6											
					0	*	*											

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 21C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Trends in Anemia by Age
Children Aged < 5 Years (2)

Page: 1
 Run Date: 2 / 2 / 07

Summary of Trends in Anemia by Age (3)							
Age and Year		Anemia Low Hb/Hct (4)		Age and Year		Anemia Low Hb/Hct (4)	
		Number	%			Number	%
6 - 11 Months				18 - 23 Months			
	2005	2,566	10.4		2005	4,522	10.3
	2004	2,775	11.1		2004	4,612	9.9
	2003	2,218	15.2		2003	4,578	10.1
	2002	59	*		2002	70	*
	2001	6,148	15.8		2001	4,603	9.8
	2000	125	14.4		2000	48	*
	1999	1	*		1999	0	*
	1998	6,522	16.1		1998	4,810	9.4
	1997	6,595	15.6		1997	4,997	10.6
12 - 17 Months				24 - 35 Months			
	2005	6,996	9.6		2005	8,751	9.2
	2004	7,179	11.1		2004	9,054	10.5
	2003	6,082	11.5		2003	8,531	11.7
	2002	114	11.4		2002	79	*
	2001	6,246	10.4		2001	8,462	11.2
	2000	85	*		2000	96	*
	1999	2	*		1999	1	*
	1998	6,029	11.1		1998	8,546	11.8
	1997	6,102	10.8		1997	8,819	11.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 21C
2005 Pediatric Nutrition Surveillance (1)
Connecticut
Summary of Trends in Anemia by Age
Children Aged < 5 Years (2)

Summary of Trends in Anemia by Age (3)							
Age and Year		Anemia Low Hb/Hct (4)		Age and Year		Anemia Low Hb/Hct (4)	
		Number	%			Number	%
36 - 59 Months				Total All Age Groups			
	2005	14,197	7.2		2005	37,032	8.7
	2004	15,104	7.4		2004	38,724	9.4
	2003	14,595	7.2		2003	36,004	9.9
	2002	232	6.9		2002	554	10.1
	2001	14,211	7.9		2001	39,670	10.4
	2000	237	6.3		2000	591	10.7
	1999	1	*		1999	5	*
	1998	15,246	8.2		1998	41,153	10.8
	1997	16,194	8.5		1997	42,707	10.8
All Other/Unknown							
	2005	0	*				
	2004	0	*				
	2003	0	*				
	2002	0	*				
	2001	0	*				
	2000	0	*				
	1999	0	*				
	1998	0	*				
	1997	0	*				

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.